

AD-A262 504 ITATION PAGE

Form Approved  
OMB No. 0704-0188

(2)



Use this space to enter the title of the report, including the title for reviewing instructions, searching existing data sources, marking the collection of information, and comments regarding the burden estimate or any other aspect of this report. Use Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Pike, Washington, DC 20540. Paperwork Reduction Project (87) 4-01. A. Washington, DC 20540.

RT DATE  
02-15-933. REPORT TYPE AND DATES COVERED  
Technical 06-01-92 to 05-31-93

## 4. TITLE AND SUBTITLE

The Synthesis and Ab Initio Structure Determination of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , a Microporous Zincoborate Built Up From "Fused" 3-Ring and 5-Ring Sub-units

## 5. FUNDING NUMBERS

N00014-90-J-1159

## 6. AUTHOR(S)

William T. A. Harrison, Thurman E. Gier, and Galen D. Stucky

## 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

University of California  
Department of Chemistry  
Santa Barbara, CA 93106

## 8. PERFORMING ORGANIZATION REPORT NUMBER

T6

## 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Office of Naval Research  
Chemistry Program  
800 N. Quincy Street  
Alexandria, VA 22217

## 10. SPONSORING/MONITORING AGENCY REPORT NUMBER

## 11. SUPPLEMENTARY NOTES

Prepared for Publication in Angewandte Chemie (accepted).

## 12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release;  
distribution unlimited

## 12b. DISTRIBUTION CODE

## 13. ABSTRACT (Maximum 200 words)

The synthesis and structure of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , a new, neutral-framework, microporous zincoborate are reported. The crystal structure of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$  has been solved ab initio from laboratory X-ray powder diffraction data, and consists of a network of vertex-sharing tetrahedral  $\text{ZnO}_4$  and triangular  $\text{BO}_3$  moieties enclosing interlinked, "cigar"-shaped cages and a three-dimensional network of intersecting channels (minimum diameter  $\sim 3.5$  Å). The "fused" 3-ring/5-ring connectivity of the Zn/B/O framework is novel. Crystal data:  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ ,  $M_r = 384.37$ , rhombohedral, space group  $R3c$  (No. 167),  $a = 9.9115$  (4) Å,  $\alpha = 48.602$  (1)°,  $V = 502.61$  Å<sup>3</sup>,  $Z = 3$ ,  $R_p = 10.9\%$ ,  $R_{wp} = 14.2\%$  for 4198 powder data (Cu K $\alpha$  radiation,  $\lambda = 1.54178$  Å).

98 4 01 069

93-06807



## 14. SUBJECT TERMS

## 15. NUMBER OF PAGES

17

## 16. PRICE CODE

17. SECURITY CLASSIFICATION  
OF REPORT  
Unclassified

18. SECURITY CLASSIFICATION  
OF THIS PAGE  
Unclassified

19. SECURITY CLASSIFICATION  
OF ABSTRACT  
Unclassified

20. LIMITATION OF ABSTRACT  
UL

NSN 7540-01-280-5500

Standard Form 298 (Rev 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

20000929137

Reproduced From  
Best Available Copy

The Synthesis and Ab Initio Structure Determination  
of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , a Microporous Zincoborate  
Built Up from "Fused" 3-Ring and 5-Ring Sub-units\*\*

By William T. A. Harrison,\* Thurman E. Gier and  
Galen D. Stucky

---

[\*] Dr. W. T. A. Harrison,

Department of Chemistry, University of Houston, Houston, TX 77205-5641 (USA)

Prof. Dr. G. D. Stucky, Dr. T. E. Gier,

Department of Chemistry, University of California, Santa Barbara, CA 93106-9510 (USA)

[\*\*] The authors gratefully acknowledge the partial financial support of the Office of Naval Research and the National Science Foundation (Grant #DMR 9208511). Dr. V. I. Srdanov (UCSB) kindly carried out the PSHG measurement.

Accepted for publication by Angewandte Chemie

# ABSTRACT

The synthesis and structure of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , a new, neutral-framework, microporous zincoborate are reported. The crystal structure of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$  has been solved *ab initio* from laboratory X-ray powder diffraction data, and consists of a network of vertex-sharing tetrahedral  $\text{ZnO}_4$  and triangular  $\text{BO}_3$  moieties enclosing interlinked, "cigar"-shaped cages and a three-dimensional network of intersecting channels (minimum diameter  $\sim 3.5 \text{ \AA}$ ). The "fused" 3-ring/5-ring connectivity of the Zn/B/O framework is novel. Crystal data:  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ ,  $M_r = 384.37$ , rhombohedral, space group  $R\bar{3}c$  (No. 167),  $a = 9.9115(4) \text{ \AA}$ ,  $\alpha = 48.602(1)^\circ$ ,  $V = 502.61 \text{ \AA}^3$ ,  $Z = 3$ ,  $R_p = 10.9\%$ ,  $R_{wp} = 14.2\%$  for 4198 powder data (Cu  $K\alpha$  radiation,  $\lambda = 1.54178 \text{ \AA}$ ).

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 4

### Short Text for Table of Contents

The syntheses, structures and properties of non-aluminosilicate molecular sieves are currently of great interest. Most previous studies have concentrated on tetrahedrally-connected frameworks, but other polyhedra types may also be incorporated into microporous structures. This communication describes  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , a zincoborate built up from tetrahedral  $\text{ZnO}_4$  and triangular  $\text{BO}_3$  sub-units.

Substitution of boron for aluminum and/or silicon in zeolite molecular sieves has long been a topic of interest.<sup>[1,2,3]</sup> A rare example of complete boron substitution for both Si and Al is provided by "boralite,"  $\text{Zn}_4\text{O}(\text{BO}_2)_6$ ,<sup>[4]</sup> a direct topological analogue of the aluminosilicate sodalite-type framework, for example,  $\text{Na}_4\text{OH}(\text{AlSiO}_4)_3$ .<sup>[5]</sup> In general, however, boron atoms, usually substituting for aluminum atoms, have only been introduced into aluminosilicate molecular sieves in small quantities, although the technological implications of such doping reactions are extremely important, as in the boron/MFI zeolite.<sup>[6]</sup>

In this communication, we describe one result of our exploratory syntheses in the zincoborate structure field, complimenting the rapidly-growing family of zincophosphate and related  $(\text{Be/Zn})(\text{P/As})$  molecular sieves,<sup>[7,8]</sup> consisting of framework elements from periodic-table-groups 2, 12 and 15.  $\text{Zn}_4\text{O}(\text{BO}_3)_2$  is a novel neutral-framework material containing micro-pores, built up from "fused", vertex-sharing tetrahedral  $\text{ZnO}_4$  and triangular  $\text{BO}_3$  units. Several other zincoborate compounds are known,<sup>[9]</sup> but none are similar to the phase described herein.

The final atomic coordinates for  $\text{Zn}_4\text{O}(\text{BO}_3)_2$  are listed in Table II, with selected geometrical data in Table III.  $\text{Zn}_4\text{O}(\text{BO}_3)_2$  has a 3-dimensional structure built up from 5 asymmetric-unit atoms (1 Zn, 1 B, 3 O), forming vertex-sharing tetrahedral  $\text{ZnO}_4$  and (almost) triangular  $\text{BO}_3$  building blocks, which enclose small, elongated cavities (Figure 1) and a three-dimensional network of channels (Figure 2). These channels propagate in the crystallographic [100], [010] and [001]-directions: Their estimated diameter is 3.5 Å. Each oxygen atom is bound to three neighbors—two zinc atoms and one boron atom for O(1) and O(2), and three zinc atoms for O(3). Average geometrical parameters are as follows:  $d_{\text{av}}(\text{Zn-O}) = 1.939(3) \text{ Å}$ ,  $d_{\text{av}}(\text{B-O}) = 1.412(4) \text{ Å}$ ,  $\theta_{\text{av}}(\text{O-Zn-O}) = 109.5(2)^\circ$  and  $\theta_{\text{av}}(\text{O-B-O}) = 120.0(4)^\circ$ . The Zn/B/O connectivity in  $\text{Zn}_4\text{O}(\text{BO}_3)_2$  leads to the presence of oxygen-bridged "3-rings" (Figure 3) and "5-rings" of nodal (Zn and B) atoms surround-

ing the central cage, although the oxygen atoms in these rings are all 3-coordinate as noted above, compared to the typical 2-fold  $T-O-T'$  connectivity in tetrahedral molecular sieve structures. The atomic-connectivity in these "fused" rings is:  $\overline{[Zn - B - Zn]}$  and  $\overline{[Zn - B - Zn - B - Zn]}$ , respectively, ignoring oxygen atoms, i.e., each ring contains a Zn-O-Zn linkage, but no B-O-B links. Extending Smith's zeolite-cage nomenclature scheme<sup>[18]</sup> to this new type of connectivity would result in an 18-nodal-atom cage with the designation  $3^65^4$  (six 3-rings and four 5-rings). These cages are aligned in the crystallographic [111]-direction, and interlinked resulting in a three-dimensional microporous network.

In summary, we have prepared and characterized a new, neutral-framework, zincoborate structure containing a novel network of small channels and cavities. Typically, a powder was produced, but the crystal structure was elucidated by *ab initio* methods from laboratory X-ray data, indicating the power of current diffraction software techniques. Further synthetic work is now in progress to determine if other novel, open, structures containing triangular boron atoms as framework species can be prepared. The presence of oxygen-atom vacancies in the framework is also notable, and may facilitate substitution reactions with the Zn/B/O framework itself.

---

#### Experimental Procedure:

**Synthesis:** The hydrothermal reaction (sealed TEFLON pouch, 200 °C, 4 days, pH  $\approx$  8.5) of 3.814 g of borax ( $Na_2B_4O_7 \cdot 10H_2O$ ), 3.440 g of 4 M NaOH solution, 6.43 g of  $Zn(NO_3)_2$  and 10 cm<sup>3</sup> of water resulted in a highly-crystalline white powder, which could be indexed "by hand" on a rhombohedral basis, with approximate lattice constants of  $a = 9.910 \text{ \AA}$  and  $\alpha = 48.62^\circ$  (hexagonal setting:  $a \approx 8.160 \text{ \AA}$ ,  $c \approx 26.156 \text{ \AA}$ ). The systematic absences of the powder data (rhombohedral setting:  $hhl$ ,  $l = 2n$ ;  $hhh$ ,  $h = 2n$ ) indicated space groups  $R3c$

or  $R\bar{3}c$ , and no "extra" lines were left un-indexed. A powder-second-harmonic-generation (PSHG) test<sup>[10]</sup> gave a null response, indicating that the material probably crystallizes in a centrosymmetric space group. Space group  $R\bar{3}c$  (No. 167) was hereafter assumed for this material, and confirmed by the successful course of the structure refinement. Thermogravimetric analysis indicated no weight loss for this material below 600 °C.

*Structure Determination:* The crystal structure of  $Zn_4O(BO_3)_2$  was solved *ab initio* from laboratory X-ray powder data.<sup>[11,12,13]</sup> High-resolution data were collected for a flat-plate sample of  $Zn_4O(BO_3)_2$  over a 12 hour period on a Scintag PAD-X automated diffractometer operating in  $\theta$ - $\theta$  geometry between  $2\theta = 16$  and  $100^\circ$ , with a step-size of  $0.02^\circ$ . Optimal estimates of the individual reflection intensities were extracted by a profile-fitting method invented by Pawley<sup>[14]</sup> and developed by Le Bail.<sup>[15]</sup> The appropriate "profile" parameters were refined by least-squares fitting and the intensities themselves (with respect to the Cu  $K\alpha_1$  wavelength) were optimized by an "iterative" procedure<sup>[15]</sup> using the program GSAS.<sup>[16]</sup> A total of 186 structure-factor-moduli ( $F_{hkl} > 0$ ), corrected for Lorentz, polarization, and multiplicity effects were obtained from the data, compared to a maximum possible number of  $\sim 230$   $F_{hkl}$  values observable to that  $(\sin\theta)/\lambda$  limit. These  $F_{hkl}$ -values were used to develop a direct methods solution (software: SHELXS-86;<sup>[17]</sup> 335 triplet-phase-relations, 7 negative-quartet-phase-relations generated) for the crystal structure of  $Zn_4O(BO_3)_2$ . Careful examination of the direct methods results indicated a plausible tetrahedral fragment [eventually atoms Zn(1) and O(1)], and the model was developed further by geometrical placement of the 2 remaining tetrahedral vertices about the center of this fragment, assuming reasonable Zn-O bond distance/angle parameters. These two  $ZnO_4$  vertices [O(2) and O(3)] were both evidently on crystallographic special positions (see Table II) and were constrained as such. These four atomic positions were used as the starting model for a conventional Rietveld refinement against the X-ray powder data.

The Rietveld refinement (program: GSAS) progressed smoothly and profile and atomic parameters were progressively added to the model as variables in the normal way. A 5-parameter pseudo-Voigt function was used to describe the X-ray line-shape and its width variation with scattering angle, and a correction for preferred orientation (direction  $\langle 111 \rangle$ ) was allowed to vary during the refinement. At the end of the initial refinement, a difference Fourier synthesis identified one distinct peak at  $\sim (0.74, 0.75, \frac{1}{4})$  which was a chemically-reasonable site for a boron atom to occupy. Bond-distance restraints were applied to this B-atom site ( $d(\text{B-O}) = 1.43(2) \text{ \AA}$ ) to assure a stable convergence of the boron-atom positional parameters. Assuming full occupancy for all five atom sites results in a unit-cell stoichiometry of  $\text{Zn}_{12}\text{B}_6\text{O}_{22}$  which does not charge-balance (apparent charge = -2). Because no extra-framework sites were apparent from Fourier maps, the oxygen fractional occupancy factors were refined. Those for O(1) and O(3) refined to values near to unity (1.00(2) and 1.06(3) respectively), while that of O(2) refined to a value of 0.84(2). The fractional occupancy of O(2) was then fixed at 0.833, and those for O(1) and O(3) were set to 1.0, giving a charge-neutral unit-cell content of  $\text{Zn}_{12}\text{B}_6\text{O}_{21}$ , or  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ . Refining this model resulted in a satisfactory convergence to final agreement values of  $R_p = 10.9\%$  and  $R_{wp} = 14.2\%$ , as defined in Table I. Further details of the crystal structure investigation may be obtained from the Fachinformationszentrum Karlsruhe, Gesellschaft für wissenschaftlich-technische Information mbH, D-W-7514 Eggenstein-Leopoldshafen 2 (FRG) on quoting the depository number CSD-\_\_\_\_, the names of the authors and the journal citation.



## References

- [1] R. Szostak, Molecular Sieves. Principles of Synthesis and Identification. Van Nostrand, New York, 1989.
- [2] D. W. Breck, Zeolite Molecular Sieves. Wiley-Interscience, New York, 1974.
- [3] M. Taramasso, G. Perego, B. Notari, in Proc. 5th Intern. Conf. Zeolites (Ed. L. V. Rees), Heyden, Philadelphia, 1980, p 205.
- [4] P. Smith-Verdier, S. Garcia-Blanco, Zeit. Krist. 1980, 151, 175.
- [5] S. Luger, J. Felsche, P. Fischer, Acta Crystallogr. 1987, C43, 1.
- [6] D. S. Santilli, S. I. Zones, in Synthesis of Microporous Materials: Vol. 1: Molecular Sieves (Eds. M. L. Occelli and H. E. Robson), Van Nostrand Reinhard, New York, 1992, p 373.
- [7] G. Harvey, W. M. Meier, in Zeolites: Facts, Figures, Future: Studies in Surface Science and Catalysis, Vol 49. (Eds. P. A. Jacobs and R. A. van Santen) Elsevier, Amsterdam, 1989, p 411.
- [8] T. F. Gier, G. D. Stucky, Nature (London) 1991, 349, 508; W. T. A. Harrison, T. E. Gier, K. L. Moran, J. M. Nicol, H. Eckert, G. D. Stucky, Chem. Mater. 1991, 3 27; T. M. Nenoff, W. T. A. Harrison, T. E. Gier, G. D. Stucky, J. Amer. Chem. Soc. 1991, 113, 378; W. T. A. Harrison, T. E. Gier, G. D. Stucky, J. Mater. Chem. 1991, 1, 153; T. E. Gier, W. T. A. Harrison, G. D. Stucky, Angew. Chem. 1991, 103, 1191; Angew. Chem. (Int. Ed. Engl.) 1991, 103, 1169. W. T. A. Harrison, T. E. Gier, T. E. Martin, G. D. Stucky. J. Mater. Chem. 1992, 2, 175; W. T. A. Harrison, T. M. Nenoff, M. M. Eddy, T.

E. Martin, G. D. Stucky. J. Mater. Chem. 1992, 2, 1127. W. T. A. Harrison, T. E. Gier, G. D. Stucky. Zeolites 1993, in press.

[9] Seven different zinc-boron-oxygen phases are included in the JCPDS powder diffraction file. Card numbers: 9-107, 9-153, 24-1438, 27-983, 27-1487, 37-1486 and 39-1126.

[10] J. P. Dougherty, S. K. Kurtz, J. Appl. Crystallogr. 1976, 9, 145.

[11] A. Le Bail, G. Ferey, P. Amoros, D. Beltran-Porter, Eur. J. Sol. St. Inorg. Chem. 1989, 26, 419.

[12] M. A. Lafontaine, A. Le Bail, G. Ferey, J. Sol. St. Chem. 1990, 85, 220.

[13] A. K. Cheetham, A. P. Wilkinson, J. Phys. Chem. Solids 1991, 52, 1199.

[14] G. S. Pawley, J. Appl. Cryst. 1981, 14, 357.

[15] A. Le Bail, H. Duroy, J. L. Fourquet, Mater. Res. Bull. 1988, 23, 447.

[16] A. C. Larson, R. B. Von Dreele, R. B. GSAS User Guide, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA, 1985-1991.

[17] G. M. Sheldrick, SHELXS-86 User Guide, University of Göttingen, Germany, 1986.

[18] J. V. Smith, Chem. Rev. 1988, 88, 149.


### Figure Captions

Fig. 1. ORTEP plot of the cage configuration in  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , viewed perpendicular to the [111]-cage axis, showing the fused 3- and 5-ring linkages: B represented by small plain circles, Zn by medium "filled" circles, O by large plain circles. Atom radii are arbitrary.

Fig. 2. ORTEP plot of the unit-cell of  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , viewed down the [100]-direction, showing the one-dimensional channels. B represented by small plain circles, Zn by medium "filled" circles, O by large plain circles. Atom radii are arbitrary.

Fig. 3. Detail of the 3-ring linkage in  $\text{Zn}_4\text{O}(\text{BO}_3)_2$ , built up from tetrahedral  $\text{ZnO}_4$  and triangular  $\text{BO}_3$  groups. Selected bond lengths [ $\text{\AA}$ ], with e.s.d.s in parentheses:  $\text{Zn}(1)-\text{O}(1)$  1.906(5),  $\text{Zn}(1)-\text{O}(1)'$  1.926(5),  $\text{Zn}(1)-\text{O}(2)$  1.962(5),  $\text{B}(1)-\text{O}(1)$  1.435(5),  $\text{B}(1)-\text{O}(2)$  1.367(7).


Table I: Crystallographic Parameters



empirical formula	$\text{Zn}_4\text{O}_7\text{B}_2$
formula wt.	384.37
crystal form	white powder
crystal system	rhombohedral
$a = b = c$ (Å)	9.9115 (4)
$\alpha = \beta = \gamma$ (°)	48.602 (1)
$V$ (Å <sup>3</sup> )	502.61
$Z$	3
space group	$R\bar{3}c$ (No. 167)
$T$ (°C)	25 (1)
$\lambda(\text{Cu K}\alpha)$ (Å)	1.54178
$\rho_{\text{calc}}$ (g/cm <sup>3</sup> )	3.81
refinement method	X-ray Rietveld refinement
software	GSAS suite
extinction correction	none applied
absorption correction	none applied
data range	$16 \leq 2\theta \leq 100^\circ$
number of data	4198 powder data points
parameters (reflections)	31 (191)
$R_p(\%)^a$	10.9
$R_{wp}(\%)^b$	14.2

<sup>a</sup> $R_p = \Sigma |y_o - C y_c| / \Sigma |y_o|$ , <sup>b</sup> $R_{wp} = [\Sigma w(y_o - C y_c)^2 / \Sigma w y_o^2]^{1/2}$ , where  $C$  is a scale factor.

Table II: Structural Data

Rhombohedral,  $R\bar{3}c$ ,  $a = 9.9115(4) \text{ \AA}$ ,  $\alpha = 48.602(1)^\circ$ 

Atom	$W^*$	$x$	$y$	$z$	$U_{iso}(\text{\AA}^2)$
Zn(1)	12f	0.6408(2)	0.6925(2)	0.0794(2)	0.007(1)
B(1)	6c	0.728(2)	0.772	1/4	0.005†
O(1)	12f	0.5854(8)	0.8045(7)	0.2181(8)	0.007(2)
O(2)§	6c	0.396(2)	3/4	0.104	0.018(4)
O(3)	4c	0.7986(4)	0.7986	0.7986	0.013(4)

\* $W$  Wyckoff-site notation, § fractional site occupancy = 5/6, †not refined

Table III: Selected Geometrical Data

Bond distances (Å)


Zn(1)-O(1)	1.906(5)	Zn(1)-O(1)'	1.926(5)
Zn(1)-O(2)	1.962(5)	Zn(1)-O(3)	1.962(2)
B(1)-O(1) × 2	1.435(5)	B(1)-O(2)	1.367(7)


Bond angles (°)

O(1)-Zn(1)-O(1)'	106.2(3)	O(1)-Zn(1)-O(2)	110.1(3)
O(1)-Zn(1)-O(3)	112.1(4)	O(1)'-Zn(1)-O(2)	111.2(2)
O(1)'-Zn(1)-O(3)	110.2(2)	O(2)-Zn(1)-O(3)	107.2(3)
O(1)-B(1)-O(1)'	120.1(8)	O(1)-B(1)-O(2)	120.0(4)
O(1)'-B(1)-O(2)	119.9(4)		

the "prime" symbol indicates a crystallographically-distinct atom



Fig. 1





$\text{Zn}_4\text{O}(\text{BO}_3)_2$





TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)\*  
Chemistry Division, Code 1113  
800 North Quincy Street  
Arlington, Virginia 22217-5000

Dr. James S. Murday (1)  
Chemistry Division, Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)  
Chemistry Division, Code 385  
Naval Air Weapons Center  
Weapons Division  
China Lake, CA 93555-6001

Dr. Elek Lindner (1)  
Naval Command, Control and Ocean  
Surveillance Center  
RDT&E Division  
San Diego, CA. 92152-5000

Dr. Bernard E. Douda (1)  
Crane Division  
Naval Surface Warfare Center  
Crane, Indiana 47522-5000

Dr. Richard W. Drisko (1)  
Naval Civil Engineering  
Laboratory  
Code L52  
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)  
Naval Surface Warfare Center  
Carderock Division Detachment  
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)  
Code 2840  
Naval Surface Warfare Center  
Carderock Division Detachment  
Annapolis, MD 21402-1198

Defense Technical Information  
Center (2)  
Building 5, Cameron Station  
Alexandria, VA 22314

\* Number of copies to forward

**END  
FILMED**

DATE:

4-93

**DTIC**